

# CHAINS OF THEORIES AND COMPANIONABILITY

ÖZCAN KASAL AND DAVID PIERCE

ABSTRACT. The theory of fields that are equipped with a countably infinite family of commuting derivations is not companionable; but if the axiom is added whereby the characteristic of the fields is zero, then the resulting theory is companionable. Each of these two theories is the union of a chain of companionable theories. In the case of characteristic zero, the model-companions of the theories in the chain form another chain, whose union is therefore the model-companion of the union of the original chain. However, in a signature with predicates, in all finite numbers of arguments, for linear dependence of vectors, the two-sorted theory of vector-spaces with their scalar-fields is companionable, and it is the union of a chain of companionable theories, but the model-companions of the theories in the chain are mutually inconsistent. Finally, the union of a chain of non-companionable theories may be companionable.

A **theory** in a given signature is a set of sentences, in the first-order logic of that signature, that is closed under logical implication. We shall consider chains  $(T_m : m \in \omega)$  of theories: this means

$$T_0 \subseteq T_1 \subseteq T_2 \subseteq \cdots \quad (*)$$

The signature of  $T_m$  will be  $\mathcal{S}_m$ , so automatically  $\mathcal{S}_0 \subseteq \mathcal{S}_1 \subseteq \mathcal{S}_2 \subseteq \cdots$

In one motivating example,  $\mathcal{S}_m$  is  $\{0, 1, -, +, \cdot, \partial_0, \dots, \partial_{m-1}\}$ , the signature of fields with  $m$  additional singularly operation-symbols; and  $T_m$  is  $m$ -DF, the theory of fields (of any characteristic) with  $m$  commuting derivations. In this example, each  $T_{m+1}$  is a **conservative extension** of  $T_m$ , that is,  $T_{m+1} \supseteq T_m$  and every sentence in  $T_{m+1}$  of signature  $\mathcal{S}_m$  is already in  $T_m$ . We establish this by showing that every model of  $T_m$  expands to a model of  $T_{m+1}$ . (This condition is sufficient, but not necessary [3, §2.6, exer. 8, p. 66].) If  $(K, \partial_0, \dots, \partial_{m-1}) \models m$ -DF, then  $(K, \partial_0, \dots, \partial_m) \models (m+1)$ -DF, where  $\partial_m$  is the 0-derivation.

---

*Date:* March 12, 2013.

*2010 Mathematics Subject Classification.* 03C10, 03C60, 12H05, 13N15.

The union of the theories  $m$ -DF can be denoted by  $\omega$ -DF: it is the theory of fields with  $\omega$ -many commuting derivations. Each of the theories  $m$ -DF has a *model-companion*, called  $m$ -DCF [11]; but we shall show (as Theorem 3 below) that  $\omega$ -DF has no model-companion. Let us recall that a **model-companion** of a theory  $T$  is a theory  $T^*$  in the same signature such that (1)  $T_{\forall} = T^*_{\forall}$ , that is, every model of one of the theories embeds in a model of the other, and (2)  $T^*$  is **model-complete**, that is,  $T^* \cup \text{diag}(\mathfrak{M})$  axiomatizes a complete theory for all models  $\mathfrak{M}$  of  $T^*$ . Here  $\text{diag}(\mathfrak{M})$  is the quantifier-free theory of  $\mathfrak{M}$  with parameters: equivalently,  $\text{diag}(\mathfrak{M})$  is the theory of all structures in which  $\mathfrak{M}_M$  embeds. (These notions, with historical references, are reviewed further in [11].) A theory has at most one model-companion, by an argument with interwoven elementary chains.

Let  $m$ -DF<sub>0</sub> be  $m$ -DF with the additional requirement that the field have characteristic 0. Then  $m$ -DF<sub>0</sub> has a model-companion, called  $m$ -DCF<sub>0</sub> [6]. We shall show (as Theorem 6 below) that  $m$ -DCF<sub>0</sub>  $\subseteq$   $(m + 1)$ -DCF<sub>0</sub>. It will follow then that the union  $\omega$ -DF<sub>0</sub> of the  $m$ -DF<sub>0</sub> has a model-companion, which is the union of the  $m$ -DCF<sub>0</sub>. This is by the following general result, which has been observed also by Alice Medvedev [7, 8]. Again, the theories  $T_k$  are as in (\*) above.

**Theorem 1.** *Suppose each theory  $T_k$  has a model-companion  $T_k^*$ , and*

$$T_0^* \subseteq T_1^* \subseteq T_2^* \subseteq \dots \quad (\dagger)$$

*Then the theory  $\bigcup_{k \in \omega} T_k$  has a model-companion, namely  $\bigcup_{k \in \omega} T_k^*$ .*

*Proof.* Write  $U$  for  $\bigcup_{k \in \omega} T_k$ , and  $U^*$  for  $\bigcup_{k \in \omega} T_k^*$ . Suppose  $\mathfrak{A} \models U$ , and  $\Gamma$  is a finite subset of  $U^* \cup \text{diag}(\mathfrak{A})$ . Then  $\Gamma$  is a subset of  $T_k^* \cup \text{diag}(\mathfrak{A} \upharpoonright \mathcal{S}_k)$  for some  $k$  in  $\omega$ , and also  $\mathfrak{A} \upharpoonright \mathcal{S}_k \models T_k$ . Since  $(T_k^*)_{\forall} \subseteq T_k$ , the structure  $\mathfrak{A} \upharpoonright \mathcal{S}_k$  must embed in a model of  $T_k^*$ ; and this model will be a model of  $\Gamma$ . We conclude that  $\Gamma$  is consistent. Therefore  $U^* \cup \text{diag}(\mathfrak{A})$  is consistent. Thus  $U^*_{\forall} \subseteq U$ . By symmetry  $U_{\forall} \subseteq U^*$ .

Similarly, if  $\mathfrak{B} \models U^*$ , then  $T_k^* \cup \text{diag}(\mathfrak{B} \upharpoonright \mathcal{S}_k)$  axiomatizes a complete theory in each case, and therefore  $U^* \cup \text{diag}(\mathfrak{B})$  is complete.  $\square$

The foregoing proof does not require that the signatures  $\mathcal{S}_k$  form a chain, but needs only that every finite subset of  $\bigcup_{k \in \omega} \mathcal{S}_k$  be included in some  $\mathcal{S}_k$ . This is the setting for Medvedev's [8, Prop. 2.4, p. 6], which then has the same proof as the foregoing. Also in Medvedev's setting, each  $T_{k+1}^*$  is a conservative extension of  $T_k^*$ ; but only the weaker assumption  $T_k^* \subseteq T_{k+1}^*$  is needed in the proof.

Medvedev notes that many properties that the theories  $T_k$  might have are ‘local’ and are therefore preserved in  $\bigcup_{k \in \omega} T_k$ : examples are completeness, elimination of quantifiers, stability, and simplicity. In her main application,  $\mathcal{S}_n$  is the signature of fields with singularly operation-symbols  $\sigma_{m/n!}$ , where  $m \in \mathbb{Z}$ ; and  $T_n$  is the theory of fields on which the  $\sigma_{m/n!}$  are automorphisms such that

$$\sigma_{k/n!} \circ \sigma_{m/n!} = \sigma_{(k+m)/n!}.$$

Then  $T_n$  includes the theory  $S_n$  of fields with the single automorphism  $\sigma_{1/n!}$ . Using [12, §1] (which is based on [3, ch. 5]), we may observe at this point that reduction of models of  $T_n$  to models of  $S_n$  is actually an equivalence of the categories  $\text{Mod}^{\subseteq}(T_n)$  and  $\text{Mod}^{\subseteq}(S_n)$ , whose objects are models of the indicated theories, and whose morphisms are embeddings. We thus have at hand a (rather simple) instance of the hypothesis of the following theorem.

**Theorem 2.** *Suppose  $(I, J)$  is a bi-interpretation of theories  $S$  and  $T$  such that  $I$  is an equivalence of the categories  $\text{Mod}^{\subseteq}(S)$  and  $\text{Mod}^{\subseteq}(T)$ . If  $S$  has the model-companion  $S^*$ , and  $S \subseteq S^*$ , then  $T$  also has a model-companion, which is the theory of those models  $\mathfrak{B}$  of  $T$  such that  $J(\mathfrak{B}) \models S^*$ .*

*Proof.* The class of models  $\mathfrak{B}$  of  $T$  such that  $J(\mathfrak{B}) \models S^*$  is elementary. Let  $T^*$  be its theory. Then  $T \subseteq T^*$ . Suppose  $\mathfrak{B} \models T$ . Then  $J(\mathfrak{B}) \models S$ , so  $J(\mathfrak{B})$  embeds in a model  $\mathfrak{A}$  of  $S^*$ . Consequently  $I(J(\mathfrak{B}))$  embeds in  $I(\mathfrak{A})$ . Also  $I(\mathfrak{A}) \models T^*$ , since  $\mathfrak{A} \cong J(I(\mathfrak{A}))$ . Since also  $\mathfrak{B} \cong I(J(\mathfrak{B}))$ , we conclude that  $\mathfrak{B}$  embeds in a model of  $T^*$ . Finally,  $T^*$  is model-complete. Indeed, suppose now  $\mathfrak{B}$  and  $\mathfrak{C}$  are models of  $T^*$  such that  $\mathfrak{B} \subseteq \mathfrak{C}$ . An embedding of  $J(\mathfrak{B})$  in  $J(\mathfrak{C})$  is induced, and these structures are models of  $S^*$ , so the embedding is elementary. Therefore the induced embedding of  $I(J(\mathfrak{B}))$  in  $I(J(\mathfrak{C}))$  is also elementary. By the equivalence of the categories,  $\mathfrak{B} \preceq \mathfrak{C}$ .  $\square$

In the present situation, the theory  $S_n$  has a model-companion [5, 1]; let us denote this by  $\text{ACFA}_n$ . By the theorem then,  $T_n$  has a model-companion  $T_n^*$ , which is axiomatized by  $T_n \cup \text{ACFA}_n$ . We have  $\text{ACFA}_n \subseteq T_{n+1}^*$  by [1, 1.12, Cor. 1, p. 3013]. By Theorem 1 then,  $\bigcup_{n \in \omega} T_n$  has a model-companion, which is the union of the  $T_n^*$ . Medvedev calls this union  $\text{QACFA}$ ; she shows for example that it preserves the simplicity of the  $\text{ACFA}_n$ , as noted above, though it does not preserve their supersimplicity.

The following is similar to the result that the theory of fields with a derivation *and* an automorphism (of the field-structure only) has no model-companion [10]. The obstruction lies in positive characteristics  $p$ , where all derivatives of elements with  $p$ -th roots must be 0.

**Theorem 3.** *The theory  $\omega$ -DF has no model-companion.*

*Proof.* We use that an  $\forall\exists$  theory  $T$  has a model-companion if and only if the class of its *existentially closed* models is elementary, and in this case the model-companion is the theory of this class [2]. (A model  $\mathfrak{A}$  of  $T$  is an **existentially closed** model, provided that if  $\mathfrak{B} \models T$  and  $\mathfrak{A} \subseteq \mathfrak{B}$ , then  $\mathfrak{A} \preceq_1 \mathfrak{B}$ , that is, all quantifier-free formulas over  $A$  that are soluble in  $\mathfrak{B}$  are soluble in  $\mathfrak{A}$ .) For each  $n$  in  $\omega$ , the theory  $\omega$ -DF has an existentially closed model  $\mathfrak{A}_n$ , whose underlying field includes  $\mathbb{F}_p(\alpha)$ , where  $\alpha$  is transcendental; and in this model,

$$\partial_k \alpha = \begin{cases} 1, & \text{if } k = n, \\ 0, & \text{otherwise.} \end{cases}$$

Then  $\alpha$  has no  $p$ -th root in  $\mathfrak{A}_n$ . Therefore, in a non-principal ultraproduct of the  $\mathfrak{A}_n$ ,  $\alpha$  has no  $p$ -th root, although  $\partial_n \alpha = 0$  for all  $n$  in  $\omega$ , so that  $\alpha$  does have a  $p$ -th root in some extension. Thus the ultraproduct is not an existentially closed model of  $\omega$ -DF. Therefore the class of existentially closed models of  $\omega$ -DF is not elementary.  $\square$

It follows then by Theorem 1 that  $m$ -DCF  $\not\leq (m+1)$ -DCF for at least one  $m$ . In fact this is so for all  $m$ , since

$$m\text{-DCF} \vdash p = 0 \rightarrow \forall x \left( \bigwedge_{i < m} \partial_i x = 0 \rightarrow \exists y y^p = x \right),$$

but  $(m+1)$ -DCF does not entail this sentence, since

$$(m+1)\text{-DCF} \vdash \exists x \left( \bigwedge_{i < m} \partial_i x = 0 \wedge \partial_m x \neq 0 \right).$$

However, this observation by itself is not enough to establish the last theorem. For, by the results of [12], it is possible for each  $T_k$  to have a model-companion  $T_k^*$ , while  $\bigcup_{k \in \omega} T_k$  has a model-companion that is not  $\bigcup_{k \in \omega} T_k^*$ . We may even require  $T_{k+1}$  to be a conservative extension of  $T_k$ .

Indeed, if  $k > 0$ , then in the notation of [12],  $\text{VS}_k$  is the theory of vector-spaces with their scalar-fields in the signature  $\{+, -, \mathbf{0}, \circ, 0, 1, *, P^k\}$ , where  $\circ$  is multiplication of scalars, and  $*$  is the action of the scalar-field on the vector-space, and  $P^k$  is  $k$ -ary linear dependence. In particular,  $P^2$  may written also as  $\|$ . Then  $\text{VS}_k$  has a model-companion,  $\text{VS}_k^*$ , which is the theory of  $k$ -dimensional vector-spaces over algebraically closed fields [12, Thm 2.3]. Let  $\text{VS}_\omega = \bigcup_{1 \leq k < \omega} \text{VS}_k$ . (This was called  $\text{VS}_\infty$  in [12].) This theory has the model-companion  $\text{VS}_\omega^*$ , which is the theory of infinite-dimensional vector-spaces over algebraically closed fields [12, Thm 2.4]. In particular

$\text{VS}_\omega^*$  is not the union of the  $\text{VS}_k^*$ , because these are mutually inconsistent. We now turn this into a result about chains:

**Theorem 4.** *If  $1 \leq n < \omega$ , let  $T_n$  be the theory axiomatized by  $\text{VS}_1 \cup \dots \cup \text{VS}_n$ . Then  $T_n$  has a model-companion  $T_n^*$ , which is axiomatized by  $T_n \cup \text{VS}_n^*$ . Also  $T_{n+1}$  is a conservative extension of  $T_n$ . However, the model-companion  $\text{VS}_\omega^*$  of the union  $\text{VS}_\omega$  of the chain  $(T_n : 1 \leq n < \omega)$  is not the union of the  $T_n^*$ .*

*Proof.* Every vector-space can be considered as a model of every  $\text{VS}_k$  and hence of every  $T_k$ . In particular,  $T_{n+1}$  is a conservative extension of  $T_n$ . If the theories  $T_n^*$  are as claimed, then they are mutually inconsistent, and so  $\text{VS}_\omega^*$  is not their union. It remains to show that there are theories  $T_n^*$  as claimed. We already know this when  $n = 1$ . For the other cases, if  $1 \leq k < n$ , we define the relations  $P^k$  in models of  $\text{VS}_n$  of dimension at least  $n$ .

Let  $\text{VS}_n^m$  the theory of such models: that is,  $\text{VS}_n^m$  is axiomatized by  $\text{VS}_n$  and the requirement that the space have dimension at least  $n$ . The relation  $P^1$  is defined in models of  $\text{VS}_n^m$  (and indeed in models of  $\text{VS}_n$ ) by the quantifier-free formula  $\mathbf{x} = \mathbf{0}$ . If  $n > 2$ , then there are existential formulas that, in each model of  $\text{VS}_n^m$ , define the relation  $\parallel$  and its complement [12, §2, p. 431]. More generally, if  $1 \leq k < n - 1$ , then, using existential formulas, we can define  $P^{k+1}$  and its complement in models of  $T_k \cup \text{VS}_n^m$  or just  $\text{VS}_k \cup \text{VS}_n^m$ . Indeed,  $\neg P^{k+1} \mathbf{x}_0 \dots \mathbf{x}_k$  is equivalent to  $\exists(\mathbf{x}_{k+1}, \dots, \mathbf{x}_{n-1}) \neg P^n \mathbf{x}_0 \dots \mathbf{x}_{n-1}$ , and  $P^{k+1} \mathbf{x}_0 \dots \mathbf{x}_k$  is equivalent to

$$\exists(\mathbf{x}_{k+1}, \dots, \mathbf{x}_n) \left( P^k \mathbf{x}_1 \dots \mathbf{x}_k \vee \left( \neg P^n \mathbf{x}_1 \dots \mathbf{x}_n \wedge \bigwedge_{j=k+1}^n P^n \mathbf{x}_0 \dots \mathbf{x}_{j-1} \mathbf{x}_{j+1} \dots \mathbf{x}_n \right) \right).$$

For, in a space of dimension at least  $n$ , if  $(\mathbf{a}_0, \dots, \mathbf{a}_k)$  is linearly dependent, but  $(\mathbf{a}_1, \dots, \mathbf{a}_k)$  is not, this means precisely that  $(\mathbf{a}_1, \dots, \mathbf{a}_n)$  is independent for some  $(\mathbf{a}_{k+1}, \dots, \mathbf{a}_n)$ , but  $\mathbf{a}_0$  is a *unique* linear combination of  $(\mathbf{a}_1, \dots, \mathbf{a}_n)$ , and in fact of  $(\mathbf{a}_1, \dots, \mathbf{a}_{j-1}, \mathbf{a}_{j+1}, \dots, \mathbf{a}_n)$  whenever  $k+1 \leq j \leq n$ , and (therefore) of  $(\mathbf{a}_1, \dots, \mathbf{a}_k)$ .

By [12, Lem 1.1, 1.2], if  $1 \leq k < n - 1$ , we now have that reduction from models of  $T_{k+1} \cup \text{VS}_n^m$  to models of  $T_k \cup \text{VS}_n^m$  is an equivalence of the categories  $\text{Mod}^{\subseteq}(T_{k+1} \cup \text{VS}_n^m)$  and  $\text{Mod}^{\subseteq}(T_k \cup \text{VS}_n^m)$ . Combining these results for all  $k$ , we have that reduction from models of  $T_{n-1} \cup \text{VS}_n^m$  to models of  $\text{VS}_n^m$  is an equivalence of the categories  $\text{Mod}^{\subseteq}(T_{n-1} \cup \text{VS}_n^m)$  and  $\text{Mod}^{\subseteq}(\text{VS}_n^m)$ . Since  $\text{VS}_n \subseteq \text{VS}_n^m$  and every model of  $\text{VS}_n$  embeds in a model

of  $VS_n^m$ , the two theories have the same model-companion, namely  $VS_n^*$ . Similarly,  $T_n$  and  $T_{n-1} \cup VS_n^m$  have the same model-companion; and by Theorem 2, this is axiomatized by  $T_n \cup VS_n^*$ .  $\square$

A one-sorted version of the last theorem can be developed as follows. Let  $VS_n^r$  comprise the sentences of  $VS_n^m$  having one-sorted signature  $\{\mathbf{0}, -, +, P^n\}$  of the sort of vectors alone. It is not obvious that all models of  $VS_n^m$  can be furnished with scalar-fields to make them models of  $VS_n^r$  again; but this will be the case. By [12, Thm 1.1], it is the case when  $n = 2$ : reduction of models of  $VS_2^m$  to models of  $VS_2^r$  is an equivalence of the categories  $\text{Mod}^{\subseteq}(VS_2^m)$  and  $\text{Mod}^{\subseteq}(VS_2^r)$ . This reduction is therefore **conservative**, by the definition of [12, p. 426]. It is said further at [12, p. 431] that reduction from  $VS_n^m$  to  $VS_n^r$  is conservative when  $n > 2$ ; but the details are not spelled out. However, the claim can be established as follows. Immediately, reduction from  $VS_2 \cup VS_n^m$  to  $VS_2^r \cup VS_n^r$  is conservative. In particular, models of the latter set of sentences really are vector-spaces without their scalar-fields. It is noted in effect in the proof of Theorem 4 that reduction from  $VS_2 \cup VS_n^m$  to  $VS_n^m$  is conservative. Furthermore, in models of the latter theory, the defining of parallelism and its complement is done with existential formulas *in the signature of vectors alone*. Therefore reduction from  $VS_2^r \cup VS_n^r$  to  $VS_n^r$  is conservative. We now have the following commutative diagram of reduction-functors, three of them being conservative, that is, being equivalences of categories.

$$\begin{array}{ccc} \text{Mod}^{\subseteq}(VS_2 \cup VS_n^m) & \longrightarrow & \text{Mod}^{\subseteq}(VS_n^m) \\ \downarrow & & \downarrow \\ \text{Mod}^{\subseteq}(VS_2^r \cup VS_n^r) & \longrightarrow & \text{Mod}^{\subseteq}(VS_n^r) \end{array}$$

Therefore the remaining reduction, from  $VS_n^m$  to  $VS_n^r$ , must be conservative.

Now there is a version of Theorem 4 where  $T_n$  is axiomatized by  $VS_2^r \cup \dots \cup VS_n^r$ . Indeed, by Theorem 2,  $T_n$  has a model-companion, which is the theory (in the same signature) of  $n$ -dimensional vector-spaces over algebraically closed fields; and the union of the  $T_n$  has a model-companion, which is the theory of infinite-dimensional vector-spaces over algebraically closed fields; but this theory is not the union of the model-companions of the  $T_n$ .

The implication  $A \Rightarrow B$  in the following is used implicitly at [1, 1.12, p. 3013] to establish the result used above, that if  $(K, \sigma)$  is a model of ACFA, then so is  $(K, \sigma^m)$ , assuming  $m \geq 1$ .

**Theorem 5.** *Assuming as usual  $T_0 \subseteq T_1$ , where each  $T_k$  has signature  $\mathcal{S}_k$ , we consider the following conditions.*

A. *For every model  $\mathfrak{A}$  of  $T_1$  and model  $\mathfrak{B}$  of  $T_0$  such that*

$$\mathfrak{A} \upharpoonright \mathcal{S}_0 \subseteq \mathfrak{B}, \quad (\ddagger)$$

*there is a model  $\mathfrak{C}$  of  $T_1$  such that*

$$\mathfrak{A} \subseteq \mathfrak{C}, \quad \mathfrak{B} \subseteq \mathfrak{C} \upharpoonright \mathcal{S}_0. \quad (\S)$$

B. *The reduct to  $\mathcal{S}_0$  of every existentially closed model of  $T_1$  is an existentially closed model of  $T_0$ .*

C.  *$T_0$  has the Amalgamation Property: if one model embeds in two others, then those two in turn embed in a fourth model, compatibly with the original embeddings.*

D.  *$T_1$  is  $\forall\exists$  (so that every model embeds in an existentially closed model).*

*We have the two implications*

$$A \implies B, \quad B \ \& \ C \ \& \ D \implies A,$$

*but there is no implication among the four conditions that does not follow from these. This is true, even if  $T_1$  is required to be a conservative extension of  $T_0$ .*

*Proof.* Suppose A holds. Let  $\mathfrak{A}$  be an existentially closed model of  $T_1$ , and let  $\mathfrak{B}$  be an arbitrary model of  $T_0$  such that  $(\ddagger)$  holds. By hypothesis, there is a model  $\mathfrak{C}$  of  $T_1$  such that  $(\S)$  holds. Then  $\mathfrak{A} \preceq_1 \mathfrak{C}$ , and therefore  $\mathfrak{A} \upharpoonright \mathcal{S}_0 \preceq_1 \mathfrak{C} \upharpoonright \mathcal{S}_0$ , and *a fortiori*  $\mathfrak{A} \upharpoonright \mathcal{S}_0 \preceq_1 \mathfrak{B}$ . Therefore  $\mathfrak{A} \upharpoonright \mathcal{S}_0$  must be an existentially closed model of  $T_0$ . Thus B holds.

Suppose conversely B holds, along with C and D. Let  $\mathfrak{A} \models T_1$  and  $\mathfrak{B} \models T_0$  such that  $(\ddagger)$  holds. We establish the consistency of  $T_1 \cup \text{diag}(\mathfrak{A}) \cup \text{diag}(\mathfrak{B})$ . It is enough to show the consistency of

$$T_1 \cup \text{diag}(\mathfrak{A}) \cup \{\exists \mathbf{x} \varphi(\mathbf{x})\}, \quad (\P)$$

where  $\varphi$  is an arbitrary quantifier-free formula of  $\mathcal{S}_0(A)$  that is soluble in  $\mathfrak{B}$ . By D, there is an existentially closed model  $\mathfrak{C}$  of  $T_1$  that extends  $\mathfrak{A}$ . By B then,  $\mathfrak{C} \upharpoonright \mathcal{S}_0$  is an existentially closed model of  $T_0$  that extends  $\mathfrak{A} \upharpoonright \mathcal{S}_0$ . By C, both  $\mathfrak{B}$  and  $\mathfrak{C} \upharpoonright \mathcal{S}_0$  embed over  $\mathfrak{A} \upharpoonright \mathcal{S}_0$  in a model of  $T_0$ . In particular,  $\varphi$  will be soluble in this model. Therefore  $\varphi$  is already soluble in  $\mathfrak{C} \upharpoonright \mathcal{S}_0$  itself. Thus  $\mathfrak{C}$  is a model of  $(\P)$ . Therefore A holds.

The foregoing arguments eliminate the five possibilities marked X on the table below, where 0 means false, and 1, true. We give examples of each

	1	X	2	3	4	X	5	6	7	X	8	9	10	X	X	11
A	0	1	0	1	0	1	0	1	0	1	0	1	0	1	0	1
B	0	0	1	1	0	0	1	1	0	0	1	1	0	0	1	1
C	0	0	0	0	1	1	1	1	0	0	0	0	1	1	1	1
D	0	0	0	0	0	0	0	0	1	1	1	1	1	1	1	1

of the remaining cases, numbered according to the table. In each example,  $T_0$  will be the reduct of  $T_1$  to  $\mathcal{S}_0$ . We shall denote by  $\mathcal{S}_f$  the signature  $\{+, \cdot, -, 0, 1\}$  of fields; and by  $\mathcal{S}_{vs}$ , the signature  $\{+, -, \mathbf{0}, \circ, 0, 1, *\}$  of vector-spaces as two-sorted structures.

1. We first give an example in which none of the four lettered conditions hold. Let  $\mathcal{S}_0 = \mathcal{S}_f \cup \{a, b\}$  and  $\mathcal{S}_1 = \mathcal{S}_0 \cup \{c\}$ . Let  $T_1$  be the theory of fields of characteristic  $p$  with distinguished elements  $a$ ,  $b$ , and  $c$  such that  $\{a, c\}$  or  $\{b, c\}$  is  $p$ -independent, and if  $\{b, c\}$  is  $p$ -independent, then so is  $\{b, c, d\}$  for some  $d$ . Then  $T_0$  is the theory of fields of characteristic  $p$  in which, for some  $c$ ,  $\{a, c\}$  or  $\{b, c\}$  is  $p$ -independent, and if  $\{b, c\}$  is  $p$ -independent, then so is  $\{b, c, d\}$  for some  $d$ . The negations of the four lettered conditions are established as follows. Throughout,  $a$ ,  $b$ ,  $c$ , and  $d$  will be algebraically independent over  $\mathbb{F}_p$ .

–A. We have

$$(\mathbb{F}_p(a, b^{1/p}, c), a, b, c) \models T_1, \quad (\mathbb{F}_p(a, b^{1/p}, c^{1/p}), a, b) \models T_0,$$

but if  $(\mathbb{F}_p(a, b^{1/p}, c), a, b, c)$  is a substructure of a model  $(K, a, b, c)$  of  $T_1$ , then  $K$  cannot contain  $c^{1/p}$ .

–B.  $T_0$  has no existentially closed models, since an element of a model that is  $p$ -independent from  $a$  or  $b$  will always have a  $p$ -th root in some extension. Similarly, no model of  $T_1$  in which  $\{a, c\}$  is not  $p$ -independent is existentially closed. But  $T_1$  does have existentially closed models, which are just the separably closed fields of characteristic  $p$  with  $p$ -basis  $\{a, c\}$  and with an additional element  $b$ .

–C.  $T_0$  does not have the Amalgamation Property, since  $(\mathbb{F}_p(a, b^{1/p}, c), a, b)$  and  $(\mathbb{F}_p(a^{1/p}, b, c, d), a, b)$  are models that do not embed in the same model over the common substructure  $(\mathbb{F}_p(a, b, c), a, b)$ , which is a model of  $T_0$ .

–D.  $T_1$  is not  $\forall\exists$ , since, as we have already noted, models in which  $\{a, c\}$  is not  $p$ -independent do not embed in existentially closed models.



2. For an example of the column headed by 2 in the table, we let  $\mathcal{S}_0$  and  $\mathcal{S}_1$  be as in 1; but now  $T_1$  is the theory of fields of characteristic  $p$  with distinguished elements  $a, b$ , and  $c$  such that  $\{a, c, d\}$  or  $\{b, c, d\}$  is  $p$ -independent for some  $d$ . This ensures that  $T_1$  has no existentially closed models, so  $B$  holds vacuously; but the other three conditions still fail.

3.  $T_0$  and  $T_1$  are the same theory, so  $A$  and  $B$  hold trivially; and this theory is the theory of vector-spaces of dimension at least 2, in the signature  $\mathcal{S}_{vs}$ , so the theory neither has the Amalgamation Property, nor is  $\forall\exists$ .

4.  $T_1$  is  $DF_p$  with the additional requirement that the field have  $p$ -dimension at least 2; and  $\mathcal{S}_0 = \mathcal{S}_f$ , so  $T_0$  is the theory of fields of characteristic  $p$  with  $p$ -dimension at least 2. The latter theory has the Amalgamation Property; but the other conditions fail. Indeed, let  $(\mathbb{F}_p(a, b), D)$  be the model of  $T_1$  in which  $Da = 1$  and  $Db = 0$ : then the field  $\mathbb{F}_p(a, b)$  embeds in  $\mathbb{F}_p(a^{1/p}, b)$ , which is a model of  $T_0$ , but  $D$  does not extend to this field. Also,  $T_0$  has no existentially closed models; but  $T_1$  does, and indeed it has a model-companion, namely  $DCF_p$ . Also  $T_1$  is not  $\forall\exists$ , since  $T_0$  is not: there is a chain of models of the latter, whose union is not a model, and we can make the structures in the chain into models of  $T_1$  by adding the zero derivation.

5.  $\mathcal{S}_0 = \mathcal{S}_f$ , and  $\mathcal{S}_1 = \mathcal{S}_0 \cup \{a\}$ .  $T_1$  is the theory of fields of characteristic  $p$  with distinguished element  $a$ , which is  $p$ -independent from another element; so  $T_0$  is (as in 4) the theory of fields of characteristic  $p$  with  $p$ -dimension at least 2. Then we already have that  $C$  holds. But  $A$  fails: just let  $\mathfrak{A}$  be  $(\mathbb{F}_p(a, b), a)$ , and let  $\mathfrak{B}$  be  $\mathbb{F}_p(a^{1/p}, b)$ . Also  $T_1$  has no existentially closed models, so  $B$  holds trivially, but  $T_1$  is not  $\forall\exists$ .

6.  $T_0$  and  $T_1$  are the same, namely the theory of fields of characteristic  $p$  of positive  $p$ -dimension, in the signature of fields, so this theory has the Amalgamation Property, but is not  $\forall\exists$ .

7.  $\mathcal{S}_0 = \mathcal{S}_{vs}$ ,  $\mathcal{S}_1 = \mathcal{S}_0 \cup \{\|\mathbf{a}, \mathbf{b}\}\}$ , and  $T_1$  is axiomatized by  $VS_2 \cup \{\mathbf{a} \nparallel \mathbf{b}\}$ , so it is  $\forall\exists$ . Then  $T_0$  is the theory of vector-spaces of dimension at least 2. As in Theorem 4 above,  $T_1$  has a model-companion, namely the theory of vector-spaces over algebraically closed fields with basis  $\{\mathbf{a}, \mathbf{b}\}$ . But  $T_0$  has no existentially closed models, since for all independent vectors  $\mathbf{a}$  and  $\mathbf{b}$  in some model, the equation

$$x * \mathbf{a} + y * \mathbf{b} = \mathbf{0} \tag{||}$$

is always soluble in some extension. Thus  $B$  fails. Then  $T_0$  also does not have the Amalgamation Property, since the solutions of (||) may satisfy  $2x^2 = y^2$  in one extension, but  $3x^2 = y^2$  in another. Similarly,  $A$  fails, since

the reduct to  $\mathcal{S}_0$  of a model of  $T_1$  may embed in a model of  $T_0$  in which  $\mathbf{a}$  and  $\mathbf{b}$  are parallel.

8.  $\mathcal{S}_0 = \mathcal{S}_{\text{vs}} \cup \{\|\}, \mathcal{S}_1 = \mathcal{S}_0 \cup \{\mathbf{a}, \mathbf{b}\}$ , and  $T_1$  is axiomatized by  $\text{VS}_2$  together with

$$\forall x \forall y (x * \mathbf{a} + y * \mathbf{b} = \mathbf{0} \rightarrow 2x^2 = y^2). \quad (**)$$

Then  $T_0$  is the theory of vector-spaces such that either the dimension is at least 2, or the scalar field contains  $\sqrt{2}$ . As in 7,  $T_0$  does not have the Amalgamation Property. The theory  $T_1$  is  $\forall\exists$ . It also has the model  $(\mathbb{Q} * \mathbf{a} \oplus \mathbb{Q} * \mathbf{b}, \mathbf{a}, \mathbf{b})$ , and  $\mathbb{Q} * \mathbf{a} \oplus \mathbb{Q} * \mathbf{b}$  embeds in the model  $\mathbb{Q}(\sqrt{2}, \sqrt{3}) * \mathbf{a}$  of  $T_0$  when we let  $\mathbf{b} = \sqrt{3} * \mathbf{a}$ ; but then the latter space embeds in no space in which  $\mathbf{a}$  and  $\mathbf{b}$  are as required by (\*\*). So  $A$  fails. Finally,  $T_1$  has a model-companion, axiomatized by  $\text{VS}_2^*$  together with

$$\exists x \exists y (x * \mathbf{a} + y * \mathbf{b} = \mathbf{0} \wedge 2x^2 = y^2 \wedge x \neq 0);$$

and  $T_0$  has a model-companion, which is just  $\text{VS}_2^*$ ; so  $B$  holds.

9.  $T_0$  and  $T_1$  are both  $\text{VS}_1$ .

10.  $T_1 = \text{DF}_p$ , and  $T_0$  is the reduct to  $\mathcal{S}_f$ , namely field-theory in characteristic  $p$ .

11.  $T_0$  and  $T_1$  are both field-theory. □

Now let  $\omega\text{-DCF}_0 = \bigcup_{m \in \omega} m\text{-DCF}_0$ . We obtain a positive application of Theorem 1.

**Theorem 6.** *For all  $m$  in  $\omega$ ,*

$$m\text{-DCF}_0 \subseteq (m+1)\text{-DCF}_0.$$

*Therefore  $\omega\text{-DF}_0$  has a model-companion, which is  $\omega\text{-DCF}_0$ . This theory admits full elimination of quantifiers, is complete, and is properly stable.*

*Proof.* Suppose  $(L, \partial_0, \dots, \partial_{m-1})$  is a model of  $m\text{-DF}_0$ , and  $L$  has a subfield  $K$  that is closed under the  $\partial_i$  (where  $i < m$ ), and there is also a derivation  $\partial_m$  on  $K$  such that  $(K, \partial_0 \upharpoonright K, \dots, \partial_{m-1} \upharpoonright K, \partial_m)$  is a model of  $(m+1)\text{-DF}_0$ . We shall include  $(L, \partial_0, \dots, \partial_{m-1})$  in another model of  $m\text{-DF}_0$ , namely a model that expands to a model of  $(m+1)\text{-DF}_0$  that includes  $(K, \partial_0, \dots, \partial_m)$ . Thus condition A of Theorem 5 will hold, and therefore condition B will hold: this means  $m\text{-DCF}_0 \subseteq (m+1)\text{-DCF}_0$ . Since  $m$  is arbitrary, it will follow by Theorem 1 that  $\omega\text{-DCF}_0$  is the model-companion of  $\omega\text{-DF}_0$ .

If  $K = L$ , we are done. Suppose  $a \in L \setminus K$ . We shall define a differential field  $(K\langle a \rangle, \tilde{\partial}_0, \dots, \tilde{\partial}_m)$ , where  $a \in K\langle a \rangle$ , and for each  $i$  in  $m$ ,

$$\tilde{\partial}_i \upharpoonright K\langle a \rangle \cap L = \partial_i \upharpoonright K\langle a \rangle \cap L, \quad (\dagger\dagger)$$

and  $\tilde{\partial}_m \upharpoonright K = \partial_m$ . Then we shall be able to repeat the process, in case  $L \not\subseteq K\langle a \rangle$ : we can work with an element of  $L \setminus K\langle a \rangle$  as we did with  $a$ . Ultimately we shall obtain the desired model of  $(m+1)$ -DF<sub>0</sub> with reduct that includes  $(L, \partial_0, \dots, \partial_{m-1})$ .

Considering  $\omega^{m+1}$  as the set of  $(m+1)$ -tuples of natural numbers, we shall have

$$K\langle a \rangle = K(a^\sigma : \sigma \in \omega^{m+1}),$$

where

$$a^\sigma = \tilde{\partial}_0^{\sigma(0)} \dots \tilde{\partial}_m^{\sigma(m)} a. \quad (\ddagger\ddagger)$$

In particular then, by  $(\ddagger\ddagger)$ , we must have

$$\sigma(m) = 0 \implies a^\sigma = \partial_0^{\sigma(0)} \dots \partial_{m-1}^{\sigma(m-1)} a.$$

Using this rule, we make the definition

$$K_1 = K(a^\sigma : \sigma(m) = 0).$$

We may assume that the derivations  $\tilde{\partial}_i$  have been defined so far that

$$i < m \implies \tilde{\partial}_i \upharpoonright K_1 = \partial_i \upharpoonright K_1, \quad \tilde{\partial}_m \upharpoonright K = \partial_m \upharpoonright K. \quad (\S\S)$$

Then  $(\ddagger\ddagger)$  holds when  $\sigma(m) < 1$ .

Now suppose that, for some positive  $j$  in  $\omega$ , we have been able to define the field  $K(a^\sigma : \sigma(m) < j)$ , and for each  $i$  in  $m$ , we have been able to define  $\tilde{\partial}_i$  as a derivation on this field, and we have been able to define  $\tilde{\partial}_m$  as a derivation from  $K(a^\sigma : \sigma(m) < j-1)$  to  $K(a^\sigma : \sigma(m) < j)$ , so that  $(\S\S)$  holds, and  $(\ddagger\ddagger)$  holds when  $\sigma(m) < j$ . We want to define the  $a^\sigma$  such that  $\sigma(m) = j$ , and we want to be able to extend the derivations  $\tilde{\partial}_i$  appropriately.

If  $i < m+1$ , then, as in [11, §4.1], we let  $\mathbf{i}$  denote the characteristic function of  $\{i\}$  on  $m+1$ : that is,  $\mathbf{i}$  will be the element of  $\omega^{m+1}$  that takes the value 1 at  $i$  and 0 elsewhere. Considered as a product structure,  $\omega^{m+1}$  inherits from  $\omega$  the binary operations  $-$  and  $+$ . For each  $i$  in  $m+1$ , we have a derivation  $\tilde{\partial}_i$  from  $K(a^\sigma : (\sigma + \mathbf{i})(m) < j)$  to  $K(a^\sigma : \sigma(m) < j)$  such that  $(\S\S)$  holds, and also, if  $\sigma(m) < j$ , then

$$\sigma(i) > 0 \implies \tilde{\partial}_i a^{\sigma - \mathbf{i}} = a^\sigma. \quad (\P\P)$$

We now define the  $a^\sigma$ , where  $\sigma(m) = j$ , so that, first of all, we can extend  $\tilde{\partial}_m$  so that  $(\P\P)$  holds when  $\sigma(m) = j$  and  $i = m$ ; but we must also ensure that  $(\P\P)$  can hold also when  $\sigma(m) = j$  and  $i < m$ . To do this, we shall have to make an inductive hypothesis, which is vacuously satisfied when  $j = 1$ . We shall also proceed recursively again. More precisely, we shall refine the recursion that we are already engaged in.

We well-order the elements  $\sigma$  of  $\omega^{m+1}$  by the linear ordering  $\triangleleft$  determined by the left-lexicographic ordering of the  $(m+1)$ -tuples

$$(\sigma(m), \sigma(0) + \cdots + \sigma(m-1), \sigma(0), \sigma(1), \dots, \sigma(m-2)).$$

Then  $(\omega^{m+1}, \triangleleft)$  has the order-type of the ordinal  $\omega^2$ . This is a difference from the linear ordering defined in [11, §4.1] and elsewhere. However, for all  $\sigma$  and  $\tau$  in  $\omega^{m+1}$ , and all  $i$  in  $m+1$ , we still have

$$\sigma \triangleleft \tau \implies \sigma + \mathbf{i} \triangleleft \tau + \mathbf{i}.$$

We have assumed that, when  $\tau = (0, \dots, 0, j)$ , we have the field  $K(a^\sigma : \sigma \triangleleft \tau)$ , together with, for each  $i$  in  $m+1$ , a derivation  $\tilde{\partial}_i$  from  $K(a^\xi : \xi + \mathbf{i} \triangleleft \tau)$  to  $K(a^\xi : \xi \triangleleft \tau)$  such that (§§) holds, and also, if  $\sigma \triangleleft \tau$ , then (¶¶) holds. We have noted that we *can* have all of this when  $\tau = (0, \dots, 0, 1)$ . Suppose we have all of this for *some*  $\tau$  in  $\omega^{m+1}$  such that  $(0, \dots, 0, 1) \trianglelefteq \tau$ , that is,  $\tau(m) > 0$ . We want to define the extension  $K(a^\sigma : \sigma \trianglelefteq \tau)$  of  $K(a^\sigma : \sigma \triangleleft \tau)$  so that we can extend the  $\tilde{\partial}_i$  appropriately. For defining  $a^\tau$ , there are two cases to consider. We use the rules for derivations gathered, for example, in [10, Fact 1.1].

1. If  $a^{\tau-\mathbf{m}}$  is algebraic over  $K(a^\xi : \xi \triangleleft \tau - \mathbf{m})$ , then the derivative  $\tilde{\partial}_m a^{\tau-\mathbf{m}}$  is determined as an element of  $K(a^\xi : \xi \triangleleft \tau)$ ; we let  $a^\tau$  be this element.
2. If  $a^{\tau-\mathbf{m}}$  is not algebraic over  $K(a^\xi : \xi \triangleleft \tau - \mathbf{m})$ , then we let  $a^\tau$  be transcendental over  $L(a^\xi : \xi \triangleleft \tau)$ . We are then free to define  $\tilde{\partial}_m a^{\tau-\mathbf{m}}$  as  $a^\tau$ . (We require  $a^\tau$  to be transcendental over  $L(a^\xi : \xi \triangleleft \tau)$ , and not just over  $K(a^\xi : \xi \triangleleft \tau)$ , so that we can establish (††) later.)

We now check that, when  $i < m$  and  $\tau(i) > 0$ , we can define  $\tilde{\partial}_i a^{\tau-\mathbf{i}}$  as  $a^\tau$ . Here we make the inductive hypothesis mentioned above, namely that the foregoing two-part definition of  $a^\tau$  was already used to define  $a^{\tau-\mathbf{i}}$ . Again we consider two cases.

1. Suppose  $a^{\tau-\mathbf{i}}$  is algebraic over  $K(a^\xi : \xi \triangleleft \tau - \mathbf{i})$ . Then  $\tilde{\partial}_i a^{\tau-\mathbf{i}}$  is determined as an element of  $K(a^\xi : \xi \triangleleft \tau)$ . Thus the value of the bracket  $[\tilde{\partial}_i, \tilde{\partial}_m]$  at  $a^{\tau-\mathbf{i}-\mathbf{m}}$  is determined: indeed, we have

$$[\tilde{\partial}_i, \tilde{\partial}_m] a^{\tau-\mathbf{i}-\mathbf{m}} = \tilde{\partial}_i \tilde{\partial}_m a^{\tau-\mathbf{i}-\mathbf{m}} - \tilde{\partial}_m \tilde{\partial}_i a^{\tau-\mathbf{i}-\mathbf{m}} = \tilde{\partial}_i a^{\tau-\mathbf{i}} - a^\tau.$$

By inductive hypothesis, since  $a^{\tau-\mathbf{i}}$  is algebraic over  $K(a^\xi : \xi \triangleleft \tau - \mathbf{i})$ , also  $a^{\tau-\mathbf{i}-\mathbf{m}}$  must be algebraic over  $K(a^\xi : \xi \triangleleft \tau - \mathbf{i} - \mathbf{m})$ . Since the bracket is 0 on this field, it must be 0 at  $a^{\tau-\mathbf{i}-\mathbf{m}}$  as well [11, Lem. 4.2].

2. If  $a^{\tau-i}$  is transcendental over  $K(a^\xi: \xi \triangleleft \tau - i)$ , then, since we are given  $\tilde{\partial}_i$  as a derivation whose domain is this field, we are free to define  $\tilde{\partial}_i a^{\tau-i}$  as  $a^\tau$ .

Thus we have obtained  $K(a^\xi: \xi \triangleleft \tau)$  as desired. By induction, we obtain the differential field  $(K(a^\sigma: \sigma \in \omega^{m+1}), \tilde{\partial}_0, \dots, \tilde{\partial}_m)$  such that  $(\dagger\dagger)$  and  $(\S\S)$  hold.

It remains to check that  $(\dagger\dagger)$  holds. It is enough to show

$$K\langle a \rangle \cap L \subseteq K_1. \quad (***)$$

(We have the reverse inclusion.) Suppose  $\tau \in \omega^{m+1}$  and  $\tau(m) > 0$ . By the definition of  $a^\tau$ ,

$$a^\tau \in K(a^\sigma: \sigma \triangleleft \tau)^{\text{alg}} \implies a^\tau \in K(a^\sigma: \sigma \triangleleft \tau), \quad (\dagger\dagger\dagger)$$

$$a^\tau \notin K(a^\sigma: \sigma \triangleleft \tau)^{\text{alg}} \implies a^\tau \notin L(a^\sigma: \sigma \triangleleft \tau)^{\text{alg}}. \quad (\dagger\dagger\dagger)$$

Suppose  $b \in K\langle a \rangle \cap L$ . Since  $b \in K\langle a \rangle$ , we have, for some  $\tau$  in  $\omega^{m+1}$ , that  $b$  is a rational function over  $K_1$  of those  $a^\sigma$  such that  $\mathbf{m} \triangleleft \sigma \triangleleft \tau$ . But then, by  $(\dagger\dagger\dagger)$ , we do not need any  $a^\sigma$  that is algebraic over  $K(a^\xi: \xi \triangleleft \sigma)$ , since it actually belongs to this field. When we throw out all such  $a^\sigma$ , then, by  $(\dagger\dagger\dagger)$ , those that remain are algebraically independent over  $L$ . Thus we have

$$b \in K_1(a^{\sigma_0}, \dots, a^{\sigma_{n-1}}) \cap L$$

for some  $\sigma_j$  in  $\omega^{m+1}$  such that  $(a^{\sigma_0}, \dots, a^{\sigma_{n-1}})$  is algebraically independent over  $L$ . Therefore we may assume  $n = 0$ , and  $b \in K_1$ . Thus  $(***)$  holds, and we have the differential field  $(K\langle a \rangle, \tilde{\partial}_0, \dots, \tilde{\partial}_m)$  fully as desired.

We have to be able to repeat this construction, in case  $L \not\subseteq K\langle a \rangle$ . If  $b \in L \setminus K\langle a \rangle$ , we have to be able to construct  $K\langle a, b \rangle$ , and so on. Let  $L\langle a \rangle$  be the compositum of  $K\langle a \rangle$  and  $L$ . Since  $m\text{-DF}_0$  has the Amalgamation Property, we can extend the  $\tilde{\partial}_i$ , where  $i < m$ , to commuting derivations on the field  $L\langle a \rangle$  that extend the original  $\tilde{\partial}_i$  on  $L$ . Thus we have a model  $(L\langle a \rangle, \tilde{\partial}_0, \dots, \tilde{\partial}_{m-1})$  of  $m\text{-DF}_0$  and a model  $(K\langle a \rangle, \tilde{\partial}_0 \upharpoonright K\langle a \rangle, \dots, \tilde{\partial}_{m-1} \upharpoonright K\langle a \rangle, \tilde{\partial}_m)$  of  $(m+1)\text{-DF}_0$  that include, respectively, the models that we started with. Now we can continue as before, ultimately extending the domain of  $\tilde{\partial}_m$  to include all of  $L$ . At limit stages of this process, we take unions, which is no problem, since  $m\text{-DF}_0$  and  $(m+1)\text{-DF}_0$  are  $\forall\exists$ .

Therefore  $\omega\text{-DF}_0$  has the model-companion  $\omega\text{-DCF}_0$ . Since the  $m\text{-DCF}_0$  have the properties of quantifier-elimination, completeness, and stability [6], the observations of Medvedev noted earlier allow us to conclude that

$\omega$ -DCF<sub>0</sub> also has these properties. Although each  $m$ -DCF<sub>0</sub> is actually  $\omega$ -stable,  $\omega$ -DCF<sub>0</sub> is not even superstable, since if  $A$  is a set of constants (in the sense that all of their derivatives are 0), then as  $\sigma$  ranges over  $A^\omega$ , the sets  $\{\partial_m x = \sigma(m) : m \in \omega\}$  belong to distinct complete types.  $\square$

In the foregoing proof, we cannot use Condition *A* of Theorem 5 in the stronger form in which the structure  $\mathfrak{C}$  is required to be a mere *expansion* to  $\mathcal{S}_1$  of  $\mathfrak{B}$ :

**Theorem 7.** *If  $m > 0$ , there is a model  $\mathfrak{K}$  of  $(m+1)$ -DF<sub>0</sub> with a reduct that is included in a model  $\mathfrak{L}$  of  $m$ -DF<sub>0</sub>, while  $\mathfrak{L}$  does not expand to a model of  $(m+1)$ -DF<sub>0</sub> that includes  $\mathfrak{K}$ .*

*Proof.* We generalize the example of [4] repeated in [9, Ex. 1.2, p. 927]. Suppose  $K$  is a pure transcendental extension  $\mathbb{Q}(a^\sigma : \sigma \in \omega^{m+1})$  of  $\mathbb{Q}$ . We make this into a model of  $(m+1)$ -DF<sub>0</sub> by requiring  $\partial_i a^\sigma = a^{\sigma+i}$  in each case. Let  $L$  be the pure transcendental extension  $K(b^\tau : \tau \in \omega^{m-1})$  of  $K$ . We make this into a model of  $m$ -DF<sub>0</sub> by extending the  $\partial_i$  so that, if  $i < m-1$ , we have  $\partial_i b^\tau = b^{\tau+i}$ , while  $\partial_{m-1} b^\tau$  is the element  $a^{(\tau,0,0)}$  of  $K$ . Note that indeed if  $i < m-1$ , then

$$[\partial_i, \partial_{m-1}]b^\tau = \partial_i a^{(\tau,0,0)} - \partial_{m-1} b^{\tau+i} = 0.$$

Suppose, if possible,  $\partial_m$  extends to  $L$  as well so as to commute with the other  $\partial_i$ . Then for any  $\tau$  in  $\omega^{m-1}$  we have  $\partial_m b^\tau = f(b^\xi : \xi \in \omega^{m-1})$  for some polynomial  $f$  over  $K$ . But then, writing  $\partial_\eta f$  for the derivative of  $f$  with respect to the variable indexed by  $\eta$ , we have, as by [10, Fact 1.1(0)],

$$\begin{aligned} a^{(\tau,0,1)} &= \partial_m \partial_{m-1} b^\tau \\ &= \partial_{m-1} \partial_m b^\tau \\ &= \partial_{m-1} (f(b^\xi : \xi \in \omega^{m-1})) \\ &= \sum_{\eta \in \omega^{m-1}} \partial_\eta f(b^\xi : \xi \in \omega^{m-1}) \cdot a^{(\eta,0,0)} + f^{\partial_{m-1}}(b^\xi : \xi \in \omega^{m-1}), \end{aligned}$$

where the sum has only finitely many nonzero terms. The polynomial expression  $f^{\partial_{m-1}}(b^\xi : \xi \in \omega^{m-1})$  cannot have  $a^{(\tau,0,1)}$  as a constant term, since this is not  $\partial_{m-1} x$  for any  $x$  in  $K$ . Thus we have obtained an algebraic relation among the  $b^\sigma$  and  $a^\tau$ ; but there can be no such relation.  $\square$

Finally, the union of a chain of non-companionable theories may be companionable:

**Theorem 8.** *In the signature  $\{f\} \cup \{c_k : k \in \omega\}$ , where  $f$  is a singular operation-symbol and the  $c_k$  are constant-symbols, let  $T_0$  be axiomatized by the sentences*

$$\forall x \forall y (fx = fy \rightarrow x = y)$$

and, for each  $k$  in  $\omega$ ,

$$\forall x (f^{k+1}x \neq x), \quad \forall x (fx = c_k \rightarrow x = c_{k+1}), \quad fc_{k+2} = c_{k+1} \rightarrow fc_{k+1} = c_k.$$

For each  $n$  in  $\omega$ , let  $T_{n+1}$  be axiomatized by

$$T_n \cup \{fc_{n+1} = c_n\}.$$

Then

- (1) each  $T_n$  is universally axiomatized, and a fortiori  $\forall\exists$ , so it does have existentially closed models;
- (2) each  $T_n$  has the Amalgamation Property;
- (3) every existentially closed model of  $T_{n+1}$  is an existentially closed model of  $T_n$ ;
- (4) no  $T_n$  is companionable;
- (5)  $\bigcup_{n \in \omega} T_n$  is companionable.

*Proof.* Let  $\mathfrak{A}_m$  be the model of  $T_0$  with universe  $\omega \times \omega$  such that

$$f^{\mathfrak{A}_m}(k, \ell) = (k, \ell + 1), \quad c_k^{\mathfrak{A}_m} = \begin{cases} (k - m, 0), & \text{if } k > m, \\ (0, m - k), & \text{if } k \leq m. \end{cases}$$

Let  $\mathfrak{A}_\omega$  be the model of  $T_0$  with universe  $\mathbb{Z}$  such that

$$f^{\mathfrak{A}_\omega}k = k + 1, \quad c_k^{\mathfrak{A}_\omega} = -k.$$

Then  $\mathfrak{A}_m$  is a model of each  $T_k$  such that  $k \leq m$ ; and  $\mathfrak{A}_\omega$  is a model of each  $T_k$ . Moreover, each model of  $T_k$  consists of a copy of some  $\mathfrak{A}_\beta$  such that  $k \leq \beta \leq \omega$ , along with some (or no) disjoint copies of  $\omega$  and  $\mathbb{Z}$  in which  $f$  is interpreted as  $x \mapsto x + 1$ . Conversely, every structure of this form is a model of  $T_k$ . The  $\beta$  such that  $\mathfrak{A}_\beta$  embeds in a given model of  $T_k$  is uniquely determined by that model. Consequently  $T_k$  has the Amalgamation Property. Also, a model of  $T_k$  is an existentially closed model if and only if it includes no copies of  $\omega$  (outside the embedded  $\mathfrak{A}_\beta$ ): This establishes that every existentially closed model of  $T_{k+1}$  is an existentially closed model of  $T_k$ .

The existentially closed models of  $T_k$  are those models that omit the type  $\{\forall y fy \neq x\} \cup \{x \neq c_j : j \in \omega\}$ . In particular,  $\mathfrak{A}_m$  is an existentially closed model of  $T_k$ , if  $k \leq m$ ; but  $\mathfrak{A}_m$  is elementarily equivalent to a structure that realizes the given type. Thus  $T_k$  is not companionable.

Finally, the model-companion of  $\bigcup_{k \in \omega} T_k$  is axiomatized by this theory, together with  $\forall x \exists y fy = x$ .  $\square$

## REFERENCES

- [1] Zoé Chatzidakis and Ehud Hrushovski, *Model theory of difference fields*, Trans. Amer. Math. Soc. **351** (1999), no. 8, 2997–3071. MR 2000f:03109
- [2] Paul Eklof and Gabriel Sabbagh, *Model-completions and modules*, Ann. Math. Logic **2** (1970/1971), no. 3, 251–295. MR MR0277372 (43 #3105)
- [3] Wilfrid Hodges, *Model theory*, Encyclopedia of Mathematics and its Applications, vol. 42, Cambridge University Press, Cambridge, 1993. MR 94e:03002
- [4] Joseph Johnson, Georg M. Reinhart, and Lee A. Rubel, *Some counterexamples to separation of variables*, J. Differential Equations **121** (1995), no. 1, 42–66. MR 96g:35006
- [5] Angus Macintyre, *Generic automorphisms of fields*, Ann. Pure Appl. Logic **88** (1997), no. 2-3, 165–180, Joint AILA-KGS Model Theory Meeting (Florence, 1995). MR 99c:03046
- [6] Tracey McGrail, *The model theory of differential fields with finitely many commuting derivations*, J. Symbolic Logic **65** (2000), no. 2, 885–913. MR 2001h:03066
- [7] Alice Medvedev,  *$\mathbb{Q}ACFA$* , Talk given at Recent Developments in Model Theory, Oléron, France, <http://modeltheory2011.univ-lyon1.fr/abstracts.html>, June 2011.
- [8] ———,  *$\mathbb{Q}ACFA$* , preprint, <http://math.berkeley.edu/~alice/grouplessqacfa.pdf>, November 2012.
- [9] David Pierce, *Differential forms in the model theory of differential fields*, J. Symbolic Logic **68** (2003), no. 3, 923–945. MR 2 000 487
- [10] ———, *Geometric characterizations of existentially closed fields with operators*, Illinois J. Math. **48** (2004), no. 4, 1321–1343. MR MR2114160
- [11] ———, *Fields with several commuting derivations*, ArXiv e-prints (2007).
- [12] ———, *Model-theory of vector-spaces over unspecified fields*, Arch. Math. Logic **48** (2009), no. 5, 421–436. MR MR2505433

MIDDLE EAST TECHNICAL UNIVERSITY, NORTHERN CYPRUS CAMPUS

MIMAR SINAN FINE ARTS UNIVERSITY, ISTANBUL

*E-mail address:* [dpierce@msgsu.edu.tr](mailto:dpierce@msgsu.edu.tr)